

# 1 The Relational Calculus

## 1.1 Basic structure

$$(R \cdot S) \cdot T = R \cdot (S \cdot T) \quad (1)$$

$$R = R \cdot id = id \cdot R \quad (2)$$

$$R \subseteq R \quad (3)$$

$$R \subseteq S \wedge S \subseteq T \Rightarrow R \subseteq T \quad (4)$$

$$R \subseteq S \wedge S \subseteq R \Rightarrow R = S \quad (5)$$

$$S \subseteq T \wedge R \subseteq U \Rightarrow S \cdot R \subseteq T \cdot U \quad (6)$$

## 1.2 Relational taxonomy

|                     |                  |             |           |
|---------------------|------------------|-------------|-----------|
| $R$ reflexive:      | $id$             | $\subseteq$ | $R$       |
| $R$ coreflexive:    | $R$              | $\subseteq$ | $id$      |
| $R$ transitive:     | $R \cdot R$      | $\subseteq$ | $R$       |
| $R$ symmetric:      | $R$              | $\subseteq$ | $R^\circ$ |
| $R$ anti-symmetric: | $R \cap R^\circ$ | $\subseteq$ | $id$      |
| $R$ connected:      | $R \cup R^\circ$ | $\subseteq$ | $\top$    |
| $R$ entire:         | $id$             | $\subseteq$ | $\ker R$  |
| $R$ simple:         | $\text{img } R$  | $\subseteq$ | $id$      |
| $R$ surjective:     | $R^\circ$        |             | entire    |
| $R$ injective:      | $R^\circ$        |             | simple    |

## 1.3 PF-transformation rules

“Guardanapo”:

$$b(f^\circ \cdot R \cdot g)a \equiv (f b)R(g a) \quad (7)$$

Left-division:

$$b(R \setminus Y)a \equiv \langle \forall c : c R b : c Y a \rangle \quad (8)$$

Pointwise ordering on functions:

$$f \sqsubseteq g \equiv f \subseteq \cdot g \equiv \langle \forall a :: (f a) \subseteq (g a) \rangle \quad (9)$$

## 1.4 Table of useful Galois connections

| $(f X) \subseteq Y \equiv X \subseteq (g Y)$ |   |  |  |
|--|---|--|--|
| Description                                  | $f = g^\flat$   | $g = f^\sharp$                             | Obs.   |
| converse                                     | $(\cdot)^\circ$   | $(\cdot)^\circ$                            |  |
| shunting 1                                   | $(h \cdot)$   | $(h^\circ \cdot)$                          | NB: $h$ is a function  |
| shunting 2                                   | $(\cdot h^\circ)$   | $(\cdot h)$                                | NB: $h$ is a function  |
| ldivision                                    | $(R \cdot)$   | $(R \setminus \cdot)$                      | $R$ under ...  |
| rdivision                                    | $(\cdot R)$   | $(\cdot / R)$                              | ... over $R$   |
| division                                     | $(R /)$   | $(\cdot \setminus R)$                      |  |
| range  | $\rho$  | $(\cdot \top)$                             | lower $\subseteq$ restricted to coreflexives                                     |
| domain                                       | $\delta$  | $(\top \cdot)$                             | lower $\subseteq$ restricted to coreflexives                                     |
| implication                                  | $(R \cap)$  | $(R \Rightarrow \cdot)$                    | Note that $(R \Rightarrow) = (\neg R \cup)$                                      |
| difference                                   | $(\cdot - R)$   | $(R \cup \cdot)$                           |  |
| PROPERTIES                                   |   |  |  |
| cancellation                                 | $X \subseteq (g \cdot f)X \quad (f \cdot g)Y \subseteq Y$ |  |  |
| definition                                   | $f X = \bigcap \{Y \mid X \subseteq gY\}$                 | $g Y = \bigcup \{X \mid f X \subseteq Y\}$ |  |
| distribution                                 | $f(X \cup Y) = (f X) \cup (f Y)$                          | $g(X \cap Y) = (g X) \cap (g Y)$           | $f(\bigcup_i X_i) = \bigcup_i (f X_i)$<br>$g(\bigcap_i X_i) = \bigcap_i (g X_i)$ |

## 1.5 Other Galois connections

Meet-universal

$$X \subseteq (R \cap S) \equiv (X \subseteq R) \wedge (X \subseteq S) \quad (10)$$

Join-universal

$$(R \cup S) \subseteq X \equiv (R \subseteq X) \wedge (S \subseteq X) \quad (11)$$

Split-universal

$$X \subseteq \langle R, S \rangle \equiv \pi_1 \cdot X \subseteq R \wedge \pi_2 \cdot X \subseteq S \quad (12)$$

Either-universal

$$X = [R, S] \equiv X \cdot i_1 = R \wedge X \cdot i_2 = S \quad (13)$$

## 1.6 “Almost” Galois connections

“Shunting” rules for  $S$  a simple relation:

$$S \cdot R \subseteq T \equiv (\delta S) \cdot R \subseteq S^\circ \cdot T \quad (14)$$

$$R \cdot S^\circ \subseteq T \equiv R \cdot \delta S \subseteq T \cdot S \quad (15)$$

Variants concerning domain and range:

$$\delta R \subseteq X \equiv R \subseteq R \cdot X \quad (16)$$

$$\rho R \subseteq X \equiv R \subseteq X \cdot R \quad (17)$$

## 1.7 Converses

$$R \subseteq R^\circ \equiv R = R^\circ \quad (18)$$

$$R \subseteq R \cdot R^\circ \cdot R \quad (19)$$

$$R^{\circ\circ} = R \quad (20)$$

$$(R \cdot S)^\circ = S^\circ \cdot R^\circ \quad (21)$$

$$(R \cap S)^\circ = R^\circ \cap S^\circ \quad (22)$$

$$(23)$$

## 1.8 Dedekind axiom

Alternative formulations: left modular law, right modular law, weak distributivity:

$$(R \cdot S) \cap T \subseteq R \cdot (S \cap (R^\circ \cdot T)) \quad (24)$$

$$(R \cdot S) \cap T \subseteq (R \cap (T \cdot S^\circ)) \cdot S \quad (25)$$

$$(R \cdot S) \cap T \subseteq (R \cap (T \cdot S^\circ)) \cdot (S \cap (R^\circ \cdot T)) \quad (26)$$

Useful instances (for  $T = id$ ,  $R = R^\circ$  and for  $S = id$ ):

$$(R^\circ \cdot S) \cap id \subseteq (R \cap S)^\circ \cdot (R \cap S) \quad (27)$$

$$R \cap T \subseteq R \cdot (id \cap (R^\circ \cdot T)) \quad (28)$$

## 1.9 Coreflexives

$$\Phi = \Phi^\circ = \Phi \cdot \Phi = \Phi \cap id \quad (29)$$

$$\Phi \cdot \Psi = \Phi \cap \Psi \quad (30)$$

$$\Phi \cdot R \subseteq S \equiv \Phi \cdot R \subseteq \Phi \cdot S \quad (31)$$

$$R \cdot \Phi \subseteq S \equiv R \cdot \Phi \subseteq S \cdot \Phi \quad (32)$$

## 1.10 Relational divisions

$$(R \setminus S) \cdot f = R \setminus (S \cdot f) \quad (33)$$

$$R \setminus (S \setminus T) = (S \cdot R) \setminus T \quad (34)$$

$$R / (S \cup T) = (R / S) \cap (R / T) \quad (35)$$

$$(S \cup T) \setminus R = (S \setminus R) \cap (T \setminus R) \quad (36)$$

## 1.11 Meets

$$R \cdot (S \cap T) = (R \cdot S) \cap (R \cdot T) \Leftarrow (\ker R) \cdot T \subseteq T \vee (\ker R) \cdot S \subseteq S \quad (37)$$

$$(S \cap T) \cdot R = (S \cdot R) \cap (T \cdot R) \Leftarrow T \cdot \text{img } R \subseteq T \vee S \cdot \text{img } R \subseteq S \quad (38)$$

$$(S \cap T) \cdot f = (S \cdot f) \cap (T \cdot f) \quad (39)$$

$$(R \cap S) \cap T \equiv R \cap (S \cap T) \quad (40)$$

$$R \cap S \equiv S \cap R \quad (41)$$

$$R \cap R \equiv R \quad (42)$$

$$R \subseteq S \equiv R = R \cap S \quad (43)$$

$$R \cap S \subseteq R \wedge R \cap S \subseteq S \quad (44)$$

$$T \cdot (R \cap S) \subseteq T \cdot R \cap T \cdot S \quad (45)$$

$$(R \cap S) \cdot T \subseteq R \cdot T \cap S \cdot T \quad (46)$$

## 1.12 Kernel and image

$$\ker R = R^\circ \cdot R \quad (47)$$

$$R \subseteq S \Rightarrow \ker R \subseteq \ker S \quad (48)$$

$$(\ker R)^\circ = \ker R \quad (49)$$

$$R \subseteq R \cdot \ker R \quad (50)$$

$$\ker (R \cdot S) = S^\circ \cdot \ker R \cdot S \quad (51)$$

$$(R^\circ \cdot S) \cap id = \ker (R \cap id) \quad (52)$$

$$\text{img } R = \ker R^\circ \quad (53)$$

### 1.13 Domain and range

$$\delta R = \ker R \cap id \quad (54)$$

$$R = R \cdot \delta R \quad (55)$$

$$\delta(R \cdot S) = \delta(\delta R \cdot S) \quad (56)$$

$$\delta(R \cap S) = (R^\circ \cdot S) \cap id \quad (57)$$

$$\rho R = \delta R^\circ \quad (58)$$

### 1.14 Splits

Definition equivalent to (12)

$$\langle R, S \rangle = \pi_1^\circ \cdot R \cap \pi_2^\circ \cdot S \quad (59)$$

The same definition pointwise: for all  $a, b, c$

$$(a, b) \langle R, S \rangle c \equiv a R c \wedge b S c \quad (60)$$

Split cancellation

$$\pi_1 \cdot \langle R, S \rangle = R \cdot \delta S \quad \wedge \quad \pi_2 \cdot \langle R, S \rangle = S \cdot \delta R \quad (61)$$

Split (conditional) fusion:

$$\langle R, S \rangle \cdot T = \langle R \cdot T, S \cdot T \rangle \Leftarrow R \cdot (\text{img } T) \subseteq R \vee S \cdot (\text{img } T) \subseteq S \quad (62)$$

Split absorption

$$\langle R \cdot T, S \cdot U \rangle = (R \times S) \cdot \langle T, U \rangle \quad (63)$$

Splits and converses:

$$\langle R, S \rangle^\circ \cdot \langle X, Y \rangle = (R^\circ \cdot X) \cap (S^\circ \cdot Y) \quad (64)$$

Therefore:

$$\ker \langle R, S \rangle = \ker R \cap \ker S \quad (65)$$

### 1.15 Eithers

Definition:

$$[R, S] = (R \cdot i_1^\circ) \cup (S \cdot i_2^\circ) \quad (66)$$

$$R + S = [i_1 \cdot R, i_2 \cdot S] \quad (67)$$

From (13), all coproduct properties extend to relations, in particular:

$$id = [i_1, i_2] \quad (68)$$

$$[R, S] \cdot i_1 = R \quad \wedge \quad [R, S] \cdot i_2 = S \quad (69)$$

$$T \cdot [R, S] = [T \cdot R, T \cdot S] \quad (70)$$

$$[T, P] \cdot (R + S) = [T \cdot R, P \cdot S] \quad (71)$$

Eithers and converses:

$$[R, S] \cdot [T, U]^\circ = (R \cdot T^\circ) \cup (S \cdot U^\circ) \quad (72)$$

$$(R + S)^\circ = R^\circ + S^\circ \quad (73)$$

### 1.16 Relational projection

Definition

$$\pi_{g,f} R \stackrel{\text{def}}{=} g \cdot R \cdot f^\circ \quad (74)$$

Property

$$\pi_{g,f} R \subseteq S \equiv g(S \leftarrow R)f \quad (75)$$

### 1.17 Equality

Ping-pong rule:

$$R = S \equiv R \subseteq S \wedge S \subseteq R \quad (76)$$

Indirection:

$$R = S \equiv \langle \forall X :: (X \subseteq R \wedge X \subseteq S) \rangle \quad (77)$$

$$R = S \equiv \langle \forall X :: (R \subseteq X \wedge S \subseteq X) \rangle \quad (78)$$